

Contemporary Management of Acute Myeloid Leukemia

A Review

Sangeetha Venugopal, MD, MS; Mikkael A. Sekeres, MD, MS

 CME at jamacmelookup.com

IMPORTANCE Acute myeloid leukemia (AML) is a clonal hematopoietic cancer that disrupts normal hematopoiesis, ultimately leading to bone marrow failure and death. The annual incidence rate of AML is 4.1 per 100 000 people in the US and is higher in patients older than 65 years. Acute myeloid leukemia includes numerous subgroups with heterogeneous molecular profiles, treatment response, and prognosis. This review discusses the evidence supporting frontline therapies in AML, the major principles that guide therapy, and progress with molecularly targeted therapy.

OBSERVATIONS Acute myeloid leukemia is a genetically complex, dynamic disease. The most commonly altered genes include *FLT3*, *NPM1*, *DNMT3A*, *IDH1*, *IDH2*, *TET2*, *RUNX1*, *NRAS*, and *TP53*. The incidence of these alterations varies by patient age, history of antecedent hematologic cancer, and previous exposure to chemotherapy and/or radiotherapy for any cancer. Since 2010, molecular data have been incorporated into AML prognostication, gradually leading to incorporation of targeted therapies into the initial treatment approach of induction chemotherapy and subsequent management. The first molecularly targeted inhibitor, midostaurin, was approved to treat patients with AML with *FLT3* variants in 2017. Since then, the understanding of the molecular pathogenesis of AML has expanded, allowing the identification of additional potential targets for drug therapy, treatment incorporation of molecularly targeted therapies (midostaurin, gilteritinib, and quizartinib targeting *FLT3* variants; ivosidenib and olutasidenib targeting *IDH1* variants, and enasidenib targeting *IDH2*), and identification of rational combination regimens. The approval of hypomethylating agents combined with venetoclax has revolutionized the therapy of AML in older adults, extending survival over monotherapy. Additionally, patients are now referred for hematopoietic cell transplant on a more rational basis.

CONCLUSIONS AND RELEVANCE In the era of genomic medicine, AML treatment is customized to the patient's comorbidities and AML genomic profile.

JAMA Oncol. doi:10.1001/jamaoncol.2024.2662
Published online August 8, 2024.

Author Affiliations: Division of Hematology, Sylvester Comprehensive Cancer Center, University of Miami, Miami, Florida.

Corresponding Author: Sangeetha Venugopal, MD, MS, Leukemia Program, Division of Hematology, Sylvester Comprehensive Cancer Center, University of Miami, 1475 NW 12th Ave, Miami, FL 33136 (svenugopal@miami.edu).

Acute myeloid leukemia (AML) is a clonal stem cell cancer characterized by proliferation of immature myeloid precursor cells and maturation arrest leading to bone marrow failure, consequent risks of infection and bleeding, and reduced overall survival (OS), with a 5-year survival rate of 31.7% (95% CI, 31.0%-32.3%).^{1,2} The annual incidence rate of AML is 4.1 per 100 000; it is more common in men (5.0 per 100 000 vs 3.4 per 100 000 for women), in non-Hispanic White individuals compared with non-Hispanic Black individuals (3.6 per 100 000), Hispanic individuals (3.4 per 100 000), and non-Hispanic Asian and Pacific Islander individuals (3.2 per 100 000), and among older adults.¹ The median age at diagnosis is 69 years, and the annual incidence rate increases to 17.3 per 100 000 in people aged 70 years and older.¹ Additional predisposing factors include previous receipt of chemotherapy³ or radiotherapy⁴ for unrelated cancers; antecedent hematologic neoplasms, such as myelodysplastic syndromes⁵ or myeloproliferative neoplasms⁶; environmental exposures to toxins, such as organic solvents⁷; and germline predisposition syndromes, such as those associated with *GATA2* or *DDX41* variants.^{8,9}

Acute myeloid leukemia has a heterogeneous cytogenomic profile, treatment susceptibility, and prognosis. Advancements in next-generation sequencing technologies and increased recognition of the molecular underpinnings of AML have led to the incorporation of molecular data in AML prognostication and treatment recommendations.¹⁰ The AML genome appears to harbor an average of 13 altered genes, of which an average of 5 are recurrently altered, including *FLT3*, *NPM1c*, *DNMT3A*, *IDH1*, *IDH2*, *TET2*, *RUNX1*, *TP53*, and *NRAS*.¹¹ Capitalizing on these advances in the understanding of AML biologic traits, since 2017, a number of AML treatments (Table) have gained regulatory approval in both molecularly defined subgroups (*FLT3* [midostaurin, quizartinib and gilteritinib], *IDH1* [ivosidenib and olutasidenib], and *IDH2* [enasidenib]) and agnostic subgroups (glasdegib, CPX-351, oral azacitidine, and combination hypomethylating agent [HMA] with venetoclax).² This has led to the incorporation of molecularly targeted agents into standard induction chemotherapy and additional therapies for patients with relapsed disease (Figure 1). Moreover, while previously older adults

Table. Commonly Used Treatments for AML

Drug	Indication	Mechanism of action	Typical dose	Efficacy data from registrational trials	Adverse effects of special interest
Gemtuzumab ozogamicin	CBF-AML	CD-33-directed antibody drug conjugate	3 mg/m ² on days 1, 4, and 7 in combination with daunorubicin and cytarabine for first induction only; for consolidation, 3 mg/m ² on day 1 in combination with cytarabine	No significant difference in CR rates between gemtuzumab ozogamicin with vs without gemtuzumab ozogamicin arms (82% vs 83%; <i>P</i> = .80); significant benefit of gemtuzumab ozogamicin observed (79% vs 51%; <i>P</i> < .001)	Grade ≥3 hemorrhages, 22.9%; veno-occlusive disease, 4.6%
Midostaurin	Newly diagnosed FLT3-altered AML, in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation	Multikinase, type 1 FLT3 inhibitor that targets both FLT3-ITD and -TKD variants	50 mg orally twice daily with food on days 8-21, with induction and consolidation	In 717 patients with FLT3-altered AML, CR rates were similar between midostaurin plus chemotherapy and PBO plus chemotherapy arms (59% vs 54%; <i>P</i> = .15); median OS, 74.6 mo in the midostaurin arm vs 25.6 mo in the PBO arm (<i>P</i> = .009)	Anemia, 93%; rash or desquamation, 14%
Quizartinib	Newly diagnosed FLT3-ITD-altered AML, in combination with chemotherapy in induction and consolidation, and as maintenance monotherapy following consolidation chemotherapy	Second-generation, type 2 FLT3 inhibitor that targets only FLT3-ITD variants	35.4 mg orally once daily on days 8-21 during induction and 35.4 mg orally once daily on days 6-19 during consolidation and continuously during maintenance therapy in those not proceeding to HCT	In 539 patients with newly diagnosed FLT3-ITD-altered AML, CR rates were similar between quizartinib plus chemotherapy and PBO plus chemotherapy arms (54.9% vs 55.4%); median OS, 31.9 mo for quizartinib plus chemotherapy vs 15.1 mo with PBO plus chemotherapy (<i>P</i> = .03)	Neutropenia, 20%; QT prolongation, 14%
CPX-351	Newly diagnosed therapy-related AML or AML with myelodysplasia-related changes	Dual drug liposomal encapsulation of cytarabine and daunorubicin that delivers a synergistic 5:1 drug ratio into leukemic blasts	Liposomal daunorubicin, 44 mg/m ² and cytarabine, 100 mg/m ² on days 1, 3, and 5 during induction vs liposomal daunorubicin, 29 mg/m ² and cytarabine, 65 mg/m ² on days 1 and 3 during consolidation	In 309 patients, CPX-351 demonstrated higher CR rate (37.3% vs 25.6%; <i>P</i> = .04) and improved OS (9.56 vs 5.95 mo; <i>P</i> = .003) vs the standard 7 + 3 regimen	Longer time to neutrophil (35 d) and platelet (36.5 d) recovery; bleeding event, 74.5%
Oral azacitidine	For continued treatment of patients with AML who achieved first CR or CRi following intensive induction chemotherapy and are unable to complete intensive curative therapy	Oral HMA administered in extended dosing schedules for sustained therapeutic activity	300 mg orally once daily on days 1-14 of each 28-d cycle	In 472 patients in first remission, azacitidine had a longer median OS vs PBO (24.7 vs 14.8 mo; <i>P</i> < .001)	Nausea, 65%; vomiting, 60%; diarrhea, 50%; neutropenia, 44%
Venetoclax with azacitidine	Newly diagnosed AML in patients aged ≥75 y or who are ineligible for IC	HMA in combination with bcl-2 inhibitor	Azacitidine, 75 mg/m ² , intravenously or subcutaneously once daily on days 1-7 of each 28-d cycle with venetoclax (dose adjusted for antifungal) daily for 28 d	In 431 patients, venetoclax with azacitidine had higher CR rates (36.7% vs 17.9%; <i>P</i> < .001) and improved median OS (14.7 vs 9.6 mo; <i>P</i> < .001) vs PBO with azacitidine	Neutropenia, 42%; febrile neutropenia, 42%; nausea, 44%; constipation, 43%; diarrhea, 41%; tumor lysis syndrome, 1%
Ivosidenib with azacitidine	Newly diagnosed IDH1-altered AML in patients aged ≥75 y or who are ineligible for IC	HMA in combination with IDH1 inhibitor	Azacitidine, 75 mg/m ² , intravenously or subcutaneously once daily on days 1-7 of each 28-d cycle with ivosidenib, 500 mg, once daily for 28 d	In 146 patients, ivosidenib with azacitidine had a CR rate of 38% vs 11% with PBO plus ivosidenib with azacitidine; median OS, 24 mo with ivosidenib plus azacitidine vs 7.9 mo with PBO plus azacitidine	Bleeding events, 41%; differentiation syndrome, 14%
Enasidenib	R/R IDH2-altered AML	IDH2 inhibitor	100 mg once daily	In 239 patients, CR rate, 20.2%; median DOR, 5.8 mo with enasidenib	Hyperbilirubinemia, 8%; differentiation syndrome, 7%

(continued)

Table. Commonly Used Treatments for AML (continued)

Drug	Indication	Mechanism of action	Typical dose	Efficacy data from registrational trials	Adverse effects of special interest
Ivosidenib	R/R <i>IDH1</i> -altered AML	<i>IDH1</i> inhibitor	500 mg once daily	In 258 patients, CR rate, 21.6%; median DOR, 9.3 mo with ivosidenib	QT prolongation, 7.8%; differentiation syndrome, 3.9%
Olutasidenib	R/R <i>IDH1</i> R132-altered AML	<i>IDH1</i> inhibitor	150 mg orally twice daily	In 153 <i>IDH1</i> inhibitor-naïve patients, CR rate, 3.2%; median DOR, 28.1 mo with olutasidenib	Nausea, 38%; differentiation syndrome, 14%; liver enzyme level increase, 12%
Gilteritinib	R/R <i>FLT3</i> -altered AML	Type 1 <i>FLT3</i> inhibitor that targets both <i>FLT3</i> -ITD and -TKD variants	120 mg orally once daily	In 371 patients, gilteritinib CR rate, 21.1%; median OS, of 9.3 mo vs salvage chemotherapy (CR, 10.5%; median OS, 5.6 mo)	Febrile neutropenia, 45.9%; anemia, 40.7%; thrombocytopenia, 22.8%; QT prolongation, 4.9%

Abbreviations: AML, acute myeloid leukemia; CBF, core-binding factor; CD, cluster differentiation; CR, complete remission; CRI, complete remission with incomplete blood count recovery; DOR, duration of response; HCT, hematopoietic cell transplant; HMA, hypomethylating agent; IC, intensive chemotherapy; ITD, internal tandem duplication; OS, overall survival; PBO, placebo; R/R, relapsed/refractory; TKD, tyrosine kinase domain.

with AML who were unable to tolerate intensive chemotherapy were treated with HMA monotherapy, if at all,¹² the approval of the HMA plus venetoclax combination has led to an incremental survival benefit and likely more older adults with AML receiving therapy.¹³ This review addresses the evidence guiding the diagnosis and treatment of adults with nonacute promyelocytic AML both in the front-line and relapsed settings.


AML Diagnosis and Prognostication

A 2-pronged approach is used in the initial evaluation of patients with AML: assessing disease-specific factors that can estimate treatment response and inform the risk of relapse (eg, prior chemotherapy or radiotherapy exposure, antecedent hematologic neoplasms and cytogenomic abnormalities) and patient-specific factors that influence an individual's ability to tolerate chemotherapy (eg, age, comorbidities, and performance status).¹⁴ Several scoring systems assess a patient's ability to tolerate therapy, incorporating factors such as age, nutritional status, preexisting organ damage, and physical function.¹⁵ Instruments measuring frailty have also correlated worse scores with impaired OS, irrespective of age.^{16,17} While there is no consensus on which scoring system best estimates an individual's ability to tolerate intensive therapy, age alone is clearly inadequate to appropriately determine up-front AML treatment.

In addition to a detailed medical history and physical examination, a bone marrow aspirate and core biopsy should be obtained to establish an AML diagnosis; the aspirate must be sent for cytogenetic analyses (karyotype and fluorescence in situ hybridization for AML-related chromosomal abnormalities) and next-generation sequencing testing for variants. In select circumstances, such as when a biopsy is technically difficult or burdensome and a patient has high levels of circulating blasts, the diagnosis can be made from peripheral blood samples. As certain variants (*FLT3*-internal tandem duplication [ITD] or *FLT3*-tyrosine kinase domain [TKD]) or specific cytogenetic abnormalities (t[8;21] or inv[16]) can immediately affect the choice of induction therapy, their assessment should be expedited.¹⁸ Central nervous system evaluation should be performed in patients with neurologic symptoms or known extramedullary leukemia.

Newly diagnosed AML is risk-stratified using the European LeukemiaNet (ELN 2022) genetic risk classification system into favorable (younger: complete remission [CR], 86% and 3-year OS rate, 75%; >60 years: CR, 81% and 3-year OS rate, 45%), intermediate (younger: CR, 59% and 3-year OS rate, 45%; >60 years: CR, 50% and 3-year OS rate, 18%), and adverse risk groups (younger: CR, 49% and 3-year OS rate, 28%; >60 years: CR, 32% and 3-year OS rate, 4%) based on relapse likelihood and survival.¹⁹ This ELN 2022 classification influences the choice of postremission therapy. Hematopoietic cell transplant (HCT) is the preferred strategy for the adverse risk group (including gene variants such as *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, or *ZRSR2*) and is considered on a case-by-case basis in the intermediate risk group (*FLT3*-ITD is intermediate risk regardless of allelic ratio or co-occurring *NPM1c*). The favorable risk group comprises core-binding factor (CBF) AML (t[8;21]: *RUNX1*-*RUNX1T1* and inv[16]: *CBFB*-*MYH11*),²⁰ altered *NPM1c* without *FLT3*-ITD,²¹ and in-frame variants affecting *CEBPA*.^{22,23} Patients with AML containing these variants receive postremission chemotherapy (Figure 2).^{24,25}

Figure 1. Mechanism of Action of Commonly Used Drugs in Acute Myeloid Leukemia (AML)



Bak, Bcl-2 antagonist killer 1; Bax, Bcl-2-associated X protein; Bcl2, B cell lymphoma 2; IDH, isocitrate dehydrogenase; ITD, internal tandem duplication; MOMP, mitochondrial outer membrane permeabilization; NADP⁺, nicotinamide adenine dinucleotide phosphate; NADPH, nicotinamide adenine dinucleotide

phosphate; Raf, rapidly accelerated fibrosarcoma; Ras, rat sarcoma; SMAC, second mitochondrial-derived activator of caspases; STAT5, signal transducer and activator of transcription; and TKD, tyrosine kinase domain; Tyk2, tyrosine kinase 2.

Management Approach in Treatment-Naïve Patients Able to Receive Intensive Induction Chemotherapy

The goal of AML therapy is to induce remission and consolidate that remission with chemotherapy and/or HCT. Younger adults (variably defined as <55-65 years) or appropriately selected older adults with a good performance status, a paucity of comorbidities, and good-risk AML are candidates for intensive chemotherapy, the backbone of which consists of cytarabine and an anthracycline (daunorubicin or idarubicin) with the addition of gemtuzumab ozogamicin in CBF-AML, midostaurin in *FLT3* (ITD or TKD)-altered AML, or quizartinib in *FLT3*-ITD-altered AML.²⁶


Core-binding factor leukemias are particularly sensitive to conventional induction and postremission chemotherapy (including intensified cytarabine), with no substantial differences in relapse-free survival or OS when cytarabine-based postremission strategies are compared with postremission HCT.²⁰ Patients with CBF-AML had improved relapse-free survival with 3 or 4 cycles of high-dose cytarabine compared with 1 or 2 cycles (7.1 vs 1.4 years; $P = .02$).²⁷ Gemtuzumab ozogamicin, a humanized anti-CD33 monoclonal antibody conjugated with the cytotoxic agent calicheamicin, can be added to standard induction therapy with daunorubicin and cytarabine for patients with newly diagnosed CBF-AML. In the UK Medical Research Council AML15 trial, patients with CBF-AML had a survival benefit with the addition of gemtuzumab ozogamicin to chemotherapy (hazard ratio [HR], 0.32; 95% CI, 0.18-0.59; $P = .001$) compared with those not receiving gemtuzumab ozogamicin.²⁸ These findings were supported in a meta-analysis of 5 randomized clinical trials (5-year OS:

76.3% with gemtuzumab ozogamicin vs 55.2% without gemtuzumab ozogamicin; HR, 0.47; 95% CI, 0.31-0.73; $P < .001$).²⁹

Assessment of measurable residual disease can potentially identify the need for additional therapy. One trial prospectively evaluated the importance of measurable residual disease by quantitative reverse transcriptase polymerase chain reaction in estimating the cumulative incidence of relapse in patients with CBF-AML. At the end of induction, a greater than 3-log reduction of *RUNX1-RUNX1T1* transcripts in the bone marrow significantly decreased relapse risk by 67% compared with a reduction of less than 3 log, while less than 10 copies of *CBFB-MYH11* in the peripheral blood was associated with a 68% reduction in relapse risk, compared with having 10 or more copies. After treatment, having more than 500 copies of *RUNX1-RUNX1T1* in bone marrow (relapse estimate: 100% vs 7% [<500 copies]; $P < .001$) and more than 10 copies of *CBFB-MYH11* with placebo (estimate of relapse: 97% vs 7% [<10 copies]; $P < .001$) were associated with relapse.³⁰ Based on these data, serial monitoring of molecular transcripts at diagnosis, remission, treatment completion, and prespecified time points during surveillance of CBF-AML is recommended to identify patients with impending relapse to consider additional therapy.

Variants of *NPM1c*-AML occurs in 20% to 30% of patients.³¹ The prognosis of *NPM1c* AML is context dependent, driven by the presence of co-occurring *FLT3*-ITD variants.²⁴ Compared with co-occurring *FLT3*-ITD variants, *NPM1c*-altered/*FLT3*-ITD wild-type AML is associated with numerically higher CR rates (74% vs 63% in *NPM1c*-altered/*FLT3*-ITD altered) and superior OS ($P = .001$) in patients treated with conventional chemotherapy.³² Measurable residual disease in *NPM1c*-altered AML should be serially monitored, as the persistence of *NPM1* transcripts at the end of induction and at treatment completion is associated with

Figure 2. Treatment Algorithm of Younger and Older Adults



AML indicates acute myeloid leukemia; CBF, core-binding factor; FISH, fluorescence in situ hybridization; HCT, hematopoietic cell therapy; IDH, isocitrate dehydrogenase; ITD, internal tandem duplication; TKD, tyrosine kinase domain; VIALE-A, venetoclax in combination with azacitidine.

increased relapse risk (HR, 4.80; 95% CI, 2.95-7.80; $P < .001$) and death (HR, 4.38; 95% CI, 2.57-7.47; $P < .001$) compared with the absence of *NPM1* transcripts.³³ The ELN 2022 guidelines recommend monitoring *NPM1* transcripts at diagnosis, at treatment completion, and 3 months for 2 years during surveillance.²⁴

Traditionally, *FLT3*-ITD variants were categorized as intermediate to adverse risk based on the allelic ratio (higher being worse).¹⁹ With the introduction of *FLT3* inhibitors into therapy, *FLT3*-ITD variant is now considered intermediate risk regardless of the allelic ratio.²⁴ The prognosis of *FLT3*-TKD alteration is variable.³⁴ Midostaurin, a small molecule, multikinase *FLT3* inhibitor, is approved to treat newly diagnosed *FLT3*-altered AML (ITD or TKD) combined with chemotherapy based on results from the RATIFY trial,³⁵ which evaluated the addition of midostaurin or placebo to induction and consolidation therapy, followed by 1 year of maintenance in patients with *FLT3*-altered AML aged 18 to 59 years. While there was no significant difference in CR rates in patients randomized to the midostaurin vs placebo arms (58.9% vs 53.5%; $P = .15$), midostaurin decreased the risk of death by 22% (HR, 0.78; 1-sided $P = .009$).³⁶

Recently quizartinib, a selective *FLT3*-ITD inhibitor, was approved to treat patients with newly diagnosed *FLT3*-ITD-altered AML based on the results of the QUANTUM-First study,³⁷ in which 539 patients with *FLT3*-ITD-altered AML were randomized to receive quizartinib ($n = 268$) or placebo ($n = 271$) combined with standard induction and postremission chemotherapy. Patients receiving quizartinib had an OS advantage compared with patients in the placebo arm (HR, 0.78; 95% CI, 0.62-0.98; $P = .03$).³⁸ Quizartinib prescribing information carries a boxed warning for QTc prolongation,

torsades de pointes, and cardiac arrest. Therefore, QTc monitoring and correction of electrolyte disturbances is recommended.³⁹ Whether quizartinib or midostaurin should be the preferred treatment in *FLT3*-ITD-altered AML is not known, as the drugs have not been compared directly.

For patients who are able to tolerate intensive chemotherapy but with ELN 2022 adverse risk genetics, the optimal treatment approach remains undefined.²⁴ With induction chemotherapy, CR rates are 48% in *TP53* variants, 42% in *KMT2A* rearrangements, and approximately 50% with monosomal karyotypes.² Whether these outcomes are better than other treatment approaches is not clear. Assessment in clinical trials is preferred, particularly for patients with *KMT2A* rearrangements who could receive targeted therapies, such as menin inhibitors.

A liposomal encapsulation of cytarabine and daunorubicin in a fixed 5:1 ratio, CPX-351 is approved to treat newly diagnosed therapy-related AML or AML with myelodysplasia-related changes in adults based on a randomized phase 3 clinical trial in patients with newly diagnosed high-risk/secondary AML ($N = 309$). Compared with a standard cytarabine- and daunorubicin-based chemotherapy regimen, CPX-351 led to a modest survival benefit (9.56 vs 5.95 months; HR, 0.69; 95% CI, 0.52-0.90; $P = .003$), including those with prior HMA exposure for antecedent myelodysplastic syndromes. The median time to neutrophil and platelet count recovery was prolonged with CPX-351 (neutrophils: 35.0 days; platelets: 36.5 days; with a 7 + 3 treatment regimen, both neutrophil and platelet count recovery was in 29 days), and use has been limited.⁴⁰

In 2020, oral azacitidine (CC-486) was approved for continued treatment of AML in patients who achieved CR or CR with

incomplete blood count recovery (CRi) following induction chemotherapy.⁴¹ QUAZAR, a randomized, placebo-controlled phase 3 trial, evaluated the role of oral azacitidine (300 mg daily in a 2 weeks on/2 weeks off schedule) or placebo in adults aged 55 years or older with AML with CR or CRi who were unable to complete standard intensive induction and postremission therapy. Patients randomized to receive oral azacitidine had an improved OS (HR, 0.69 vs placebo; 95% CI, 0.55-0.86; $P < .001$).⁴² Most importantly, oral azacitidine is not a pharmacologic equivalent of subcutaneous or intravenous azacitidine and the formulations are not interchangeable.^{43,44}

Management Approach in Treatment-Naïve Patients Unable to Receive Intensive Induction Chemotherapy

Acute myeloid leukemia is more common in older adults, who also are more likely to have concomitant comorbidities and poor performance status.⁴⁵ Among patients aged 70 years or older or younger patients with poor performance status and/or major comorbidities, intensive chemotherapy is associated with a high risk of early mortality and truncated survival.⁴⁶ Likely as a consequence, 50% of older adults with AML between 2001 and 2013 did not receive any active therapy, including 42% with AML diagnosed in the last decade.⁴⁷ The 2020 approval of venetoclax (an orally available Bcl2 inhibitor) in combination with HMAs for newly diagnosed AML in older adults (aged ≥ 75 years or unable to receive intensive chemotherapy) has expanded the AML treatment landscape for this population.⁴⁸

VIALE-A, a randomized, double-blind, placebo-controlled phase 3 study, investigated the efficacy and safety of venetoclax in combination with azacitidine compared with placebo plus azacitidine in patients with newly diagnosed AML who were older (age ≥ 75 years) or unlikely to tolerate intensive chemotherapy.¹³ The composite CR rate (CRc: CR plus CRi) in the venetoclax plus azacitidine arm (64.7%) was significantly higher than in the placebo plus azacitidine arm (22.8%) ($P < .001$). At a median follow-up of 20.5 months, venetoclax plus azacitidine decreased the risk of death by 33.8% (HR, 0.662; $P < .001$), with a median survival duration of 14.7 vs 9.6 months for patients receiving placebo plus azacitidine.¹³

Focusing on patients with discrete variants, in those with an *FLT3* variant (ITD or TKD) treated with venetoclax plus azacitidine, the CRc rate was 67% and the median OS was 12.5 months (placebo plus azacitidine: CRc, 36%; median OS, 8.6 months).⁴⁹ Among patients with *IDH* variants or *IDH2* variants treated with venetoclax plus azacitidine, the CRc was 79% and the median OS was 24.5 months (placebo plus azacitidine: CRc, 11%; median OS, 6.2 months).⁵⁰

In the subgroup analysis of patients with adverse cytogenetics, outcomes with venetoclax plus azacitidine were influenced by the presence or absence of *TP53* variation. Although the subgroup analysis was not powered to detect differences, venetoclax plus azacitidine treatment was associated with numerically higher response rates and OS (CRc, 70%; OS, 23.4 months) compared with placebo plus azacitidine (CRc, 23%; OS, 11.3 months) in those with adverse cytogenetics and *TP53* wild type. However, among patients with adverse cytogenetics and *TP53* variants (a surrogate marker for multi-hit *TP53*⁵¹), survival appeared similar between the arms (veneto-

clax plus azacitidine, 5.2 months; placebo plus azacitidine, 4.9 months).⁵² Taken together, venetoclax plus azacitidine appears to be an effective therapy across the molecular subgroups in older adults with AML and those unable to tolerate intensive chemotherapy and should be continued ad infinitum, as long as a patient is deriving benefit. Venetoclax-based regimens can also be used when measurable residual disease clearance is suboptimal.⁵³ Whether the combination truly represents nonintensive therapy, whether outcomes differ from intensive induction chemotherapy in the same or in a younger population,⁵⁴ and the role of triplet therapy in patients with *FLT3*-altered AML (HMA plus venetoclax plus *FLT3* inhibitor) is still being investigated.^{55,56} Use of checkpoint inhibitors combined with standard chemotherapy is an area of active investigation.

In 2020, ivosidenib was approved to treat patients with relapsed or refractory (R/R) *IDH1*-altered AML, and in an extended approval in 2022, ivosidenib in combination with azacitidine was approved for newly diagnosed *IDH1*-altered AML in adults aged 75 years or older or who are unable to receive intensive induction chemotherapy. AGILE, a placebo-controlled, randomized phase 3 trial, enrolled 146 patients with newly diagnosed *IDH1*-altered AML to receive ivosidenib plus azacitidine or placebo plus azacitidine, with the primary end point of event-free survival. At a median follow-up of 12.4 months, the event-free survival was significantly longer for those randomized to ivosidenib plus azacitidine compared with the placebo plus azacitidine group (HR, 0.33; 95% CI, 0.16-0.69; $P = .002$). Differentiation syndrome, an ivosidenib-specific adverse event, occurred in 14% of patients receiving ivosidenib plus azacitidine vs 9% with placebo plus azacitidine.⁵⁷ Whether older adults with *IDH1* variants or those unlikely to tolerate intensive chemotherapy should receive initial ivosidenib plus azacitidine or venetoclax plus azacitidine is being investigated.⁵⁸

Management of Relapsed AML

Despite the major headway in unraveling the byzantine molecular pathogenesis of AML and the identification of rational therapies, relapse is more common, with a 5-year OS of 10% in patients whose AML returns.⁵⁹ Currently, there is no consensus on a uniform reinduction strategy in patients who can tolerate intensive chemotherapy and who do not have actionable variants, and the choice of treatment is often based on the CR duration with the prior therapy. If the first remission duration is greater than 1 year, it is reasonable to reinduce with the same initial regimen. However, the likelihood of treatment response decreases with each relapse. Commonly used intensive chemotherapy regimens in the R/R setting include the mitoxantrone, etoposide, and cytarabine combination; fludarabine, cytarabine, granulocyte colony-stimulating factor, and idarubicin combination; or high-dose cytarabine (3 g/m²). In patients not able to receive intensive chemotherapy, other regimens (eg, hypomethylating agents combined with venetoclax in those previously treated with cytotoxic chemotherapy) may be used.⁶⁰ The efficacy of these regimens is underwhelming, with CR rates ranging from 30% to 65% and a median duration of response typically less than 12 months.²⁶ The only cure for R/R AML is HCT, which may not require treatment before the transplantation-conditioning regimen. In the phase 3 ASAP trial, pre-HCT sequential conditioning with high-dose cytarabine or melphalan followed by reduced-

intensity conditioning in emergent HCT was noninferior to remission-induction chemotherapy prior to HCT (1-year leukemia-free survival: sequential conditioning, 71.5% vs remission-induction chemotherapy, 69.9%; $P = .80$).⁶¹ Clinical trial options must be considered when available.

Patients with discrete, targetable variants have other therapeutic options. Enasidenib, an orally available, small-molecule IDH2 inhibitor, is approved to treat patients with R/R IDH2-altered AML. Its approval was based on the phase 1-2 AG-221-C-001 study, for which the primary end point was overall response rate (CR, CRi, CR with incomplete platelet recovery, partial remission, or morphologic leukemia-free state).⁶² Enasidenib demonstrated an overall response rate of 40.3% and a CR or CR with partial hematologic recovery rate of 23% lasting a median of 5.8 months. Additionally, 34% of the patients became transfusion independent (red blood cells or platelets) during any 56-day postbaseline period. Enasidenib-specific adverse events included indirect hyperbilirubinemia (12%) and differentiation syndrome (7%). In the phase 3 study comparing enasidenib with conventional care in a similar patient population, however, enasidenib did not demonstrate a survival benefit (6.5 vs 6.2 months; $P = .23$), indicating that while it is a treatment option, it may not necessarily be a preferred treatment option.⁶³

In the past 5 years, 2 IDH1 inhibitors, (ivosidenib and olutasidenib) have been approved to treat R/R IDH1-altered AML.⁶⁴ Ivosidenib and olutasidenib are both allosteric type II IDH1 inhibitors, but differ in their chemical structure and binding properties.^{65,66} In the phase 1b trial that led to the approval of ivosidenib in R/R IDH1-altered AML, patients treated with ivosidenib had an overall response rate of 41.6%, including a CR or CR with partial hematologic recovery rate of 30.6%, and a median duration of response of 8.2 months among responders.⁶⁷ In the pivotal trial that led to the approval of olutasidenib in a similar population, the overall response rate was 48%, including a CR or CR with partial hematologic recovery rate of 35% and a median duration of response of 25.9 months among responders.⁶⁸ Ivosidenib and olutasidenib differ slightly in their adverse event profile, with ivosidenib carrying a boxed warning for QTc interval prolongation and olutasidenib for hepatotoxicity.⁶⁴ Given the lack of head-to-head comparison between ivosidenib and olutasidenib, the choice of IDH1 inhibitor in R/R IDH1-altered AML depends on likely patient tolerability, comorbidities, and prior exposure to other therapies.

Gilteritinib, a highly selective, oral FLT3 inhibitor active against ITD and TKD variants, is approved in the US to treat R/R FLT3-altered AML based on results of the ADMIRAL study.⁶⁹ Gilteritinib was compared with salvage chemotherapy (mitoxantrone, etoposide, and cytarabine combination; fludarabine, cytarabine,

granulocyte colony-stimulating factor, and idarubicin combination; or low-dose cytarabine or azacitidine) in 371 patients with R/R FLT3-altered AML.⁷⁰ The CR or CR with partial hematologic recovery rates were significantly higher in patients treated with gilteritinib compared with patients treated with salvage chemotherapy (34% vs 15.3%; risk difference, 18.6%; 95% CI, 9.8%-27.4%). At a median follow-up of 17.8 months, gilteritinib demonstrated a significant improvement in OS compared with salvage chemotherapy (9.3 vs 5.6 months; HR, 0.64; 95% CI, 0.49-0.83; $P < .001$).⁷¹ The gilteritinib labeling includes a boxed warning for differentiation syndrome and warnings for posterior reversible encephalopathy syndrome and QTc prolongation.⁶⁹

Conclusions

Over the past decade, therapy for patients with AML has increasingly become tailored to both the disease molecular profile and patient- and disease-specific characteristics. Novel orally administered targeted agents pose coverage challenges and may necessitate paritylike legislative approaches to ensure patients can receive standard-of-care therapies regardless of administrative route. Drug development in AML is proceeding at a vigorous pace with continued exploitation of disease biologic factors, such as with the ongoing investigation of menin inhibitors against *KMT2A* and *NPM1c* variants.⁷² Nevertheless, relapse, treatment refractoriness (common with *TP53* multihit status), and emergence of novel resistance variants remain major impediments to successful therapy.^{73,74} Patients should be considered for clinical trials in up-front, relapsed, or refractory settings when available, given the persistent poor outcomes of this population. The approval of HMA plus venetoclax in older adults and those unlikely to tolerate intensive chemotherapy has spurred the evaluation of triplet therapies with an add-on third agent (either approved or investigational) to the HMA plus venetoclax backbone. Additional core questions, such as the optimal duration of venetoclax therapy, time-limited treatment, or appropriate sequencing of HMA plus venetoclax and molecularly targeted therapies, are under active investigation.⁷⁵⁻⁷⁷ The concept of measurable residual disease assessment in AML is gaining traction and will likely serve as a marker to guide therapy duration and need for additional treatment interventions. Given the important prognostic information imparted by the cytogenomic makeup of AML, biomarker-focused clinical research using advanced approaches, such as basket, umbrella, and adaptive study designs, will accelerate drug development timelines. To quote Mark Twain, "the secret of getting ahead is getting started,"⁷⁸ and therapeutic development intended to improve the survival outcomes in patients with AML is well on its way.

ARTICLE INFORMATION

Accepted for Publication: March 12, 2024.

Published Online: August 8, 2024.
doi:10.1001/jamaoncol.2024.2662

Author Contributions: Drs Venugopal and Sekeres had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.
Concept and design: Both authors.
Acquisition, analysis, or interpretation of data: Venugopal.

Drafting of the manuscript: Both authors.
Critical review of the manuscript for important intellectual content: Both authors.
Administrative, technical, or material support: Sekeres.
Supervision: Sekeres.

Conflict of Interest Disclosures: Dr Venugopal reported receiving personal fees from serving on the Ascentage Advisory Board. Dr Sekeres reported receiving personal fees from serving on the BMS Advisory Board, Kurome Advisory Board, and

Novartis Advisory Board outside the submitted work. No other disclosures were reported.

REFERENCES

- SEER*Explorer: an interactive website for SEER cancer statistics. Surveillance Research Program, National Cancer Institute. December 16, 2023. <https://seer.cancer.gov/statistics-network/explorer/>
- DiNardo CD, Erba HP, Freeman SD, Wei AH. Acute myeloid leukaemia. *Lancet*. 2023;401

- (10393):2073-2086. doi:10.1016/S0140-6736(23)00108-3
3. Rosenstock AS, Niu J, Giordano SH, Zhao H, Wolff AC, Chavez-MacGregor M. Acute myeloid leukemia and myelodysplastic syndrome after adjuvant chemotherapy: a population-based study among older breast cancer patients. *Cancer*. 2018;124(5):899-906. doi:10.1002/cncr.31144
 4. Schonfeldt SJ, Gilbert ES, Dores GM, et al. Acute myeloid leukemia following Hodgkin lymphoma: a population-based study of 35,511 patients. *J Natl Cancer Inst*. 2006;98(3):215-218. doi:10.1093/jnci/dji017
 5. Granfeldt Østgård LS, Medeiros BC, Sengeløv H, et al. Epidemiology and clinical significance of secondary and therapy-related acute myeloid leukemia: a national population-based cohort study. *J Clin Oncol*. 2015;33(31):3641-3649. doi:10.1200/JCO.2014.60.0890
 6. Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood*. 2014;124(16):2507-2513. doi:10.1182/blood-2014-05-579136
 7. McHale CM, Zhang L, Smith MT. Current understanding of the mechanism of benzene-induced leukemia in humans: implications for risk assessment. *Carcinogenesis*. 2012;33(2):240-252. doi:10.1093/carcin/bgr297
 8. Shallis RM, Wang R, Davidoff A, Ma X, Zeidan AM. Epidemiology of acute myeloid leukemia: recent progress and enduring challenges. *Blood Rev*. 2019;36:70-87. doi:10.1016/j.blre.2019.04.005
 9. Polprasert C, Schulze I, Sekeres MA, et al. Inherited and somatic defects in *DDX41* in myeloid neoplasms. *Cancer Cell*. 2015;27(5):658-670. doi:10.1016/j.ccell.2015.03.017
 10. Döhner H, Estey EH, Amadori S, et al; European LeukemiaNet. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453-474. doi:10.1182/blood-2009-07-235358
 11. Ley TJ, Miller C, Ding L, et al; Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368(22):2059-2074. doi:10.1056/NEJMoa1301689
 12. Zeidan AM, Wang R, Wang X, et al. Clinical outcomes of older patients with AML receiving hypomethylating agents: a large population-based study in the United States. *Blood Adv*. 2020;4(10):2192-2201. doi:10.1182/bloodadvances.2020001779
 13. DiNardo CD, Jonas BA, Pullarkat V, et al. Azacitidine and venetoclax in previously untreated acute myeloid leukemia. *N Engl J Med*. 2020;383(7):617-629. doi:10.1056/NEJMoa2012971
 14. Cortes JE, Mehta P. Determination of fitness and therapeutic options in older patients with acute myeloid leukemia. *Am J Hematol*. 2021;96(4):493-507. doi:10.1002/ajh.26079
 15. Palmieri R, Othus M, Halpern AB, et al. Accuracy of SIE/SIES/GITMO consensus criteria for unfit patients to predict early mortality after intensive chemotherapy in adults with AML or other high-grade myeloid neoplasm. *J Clin Oncol*. 2020;38(35):4163-4174. doi:10.1200/JCO.20.01392
 16. Umit EG, Baysal M, Demir AM. Frailty in patients with acute myeloid leukaemia, conceptual misapprehension of chronological age. *Eur J Cancer Care (Engl)*. 2018;27(2):e12810. doi:10.1111/ecc.12810
 17. Lee MH, La J, Brophy M, et al. Baseline correlates of frailty and its association with survival in US veterans with newly diagnosed and treated acute myeloid leukemia. *Blood*. 2022;140(suppl 1):8117-8118. doi:10.1182/blood-2022-169225
 18. Short NJ, Rytting ME, Cortes JE. Acute myeloid leukaemia. *Lancet*. 2018;392(10147):593-606. doi:10.1016/S0140-6736(18)31041-9
 19. Döhner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017;129(4):424-447. doi:10.1182/blood-2016-08-733196
 20. Schlenk RF, Benner A, Krauter J, et al. Individual patient data-based meta-analysis of patients aged 16 to 60 years with core binding factor acute myeloid leukemia: a survey of the German Acute Myeloid Leukemia Intergroup. *J Clin Oncol*. 2004;22(18):3741-3750. doi:10.1200/JCO.2004.03.012
 21. Thiede C, Koch S, Creutzig E, et al. Prevalence and prognostic impact of *NPM1* mutations in 1485 adult patients with acute myeloid leukemia (AML). *Blood*. 2006;107(10):4011-4020. doi:10.1182/blood-2005-08-3167
 22. Taube F, Georgi JA, Kramer M, et al; Study Alliance Leukemia (SAL). *CEBPA* mutations in 4708 patients with acute myeloid leukemia: differential impact of bZIP and TAD mutations on outcome. *Blood*. 2022;139(1):87-103. doi:10.1182/blood.2020009680
 23. Wakita S, Sakaguchi M, Oh I, et al. Prognostic impact of *CEBPA* bZIP domain mutation in acute myeloid leukemia. *Blood Adv*. 2022;6(1):238-247. doi:10.1182/bloodadvances.2021004292
 24. Döhner H, Wei AH, Appelbaum FR, et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022;140(12):1345-1377. doi:10.1182/blood.2022016867
 25. Short NJ, Tallman MS, Pollyea DA, Ravandi F, Kantarjian H. Optimizing risk stratification in acute myeloid leukemia: dynamic models for a dynamic therapeutic landscape. *J Clin Oncol*. 2021;39(23):2535-2538. doi:10.1200/JCO.21.00067
 26. Pollyea DA, Altman JK, Assi R, et al. Acute myeloid leukemia, version 3.2023, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw*. 2023;21(5):503-513. doi:10.6004/jncn.2023.0025
 27. Byrd JC, Ruppert AS, Mrózek K, et al. Repetitive cycles of high-dose cytarabine benefit patients with acute myeloid leukemia and inv(16)(p13q22) or t(16;16)(p13;q22): results from CALGB 8461. *J Clin Oncol*. 2004;22(6):1087-1094. doi:10.1200/JCO.2004.07.012
 28. Burnett AK, Hills RK, Milligan D, et al. Identification of patients with acute myeloblastic leukemia who benefit from the addition of gemtuzumab ozogamicin: results of the MRC AML15 trial. *J Clin Oncol*. 2011;29(4):369-377. doi:10.1200/JCO.2010.31.4310
 29. Hills RK, Castaigne S, Appelbaum FR, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy in adult patients with acute myeloid leukaemia: a meta-analysis of individual patient data from randomised controlled trials. *Lancet Oncol*. 2014;15(9):986-996. doi:10.1016/S1470-2045(14)70281-5
 30. Yin JA, O'Brien MA, Hills RK, Daly SB, Wheatley K, Burnett AK. Minimal residual disease monitoring by quantitative RT-PCR in core binding factor AML allows risk stratification and predicts relapse: results of the United Kingdom MRC AML15 trial. *Blood*. 2012;120(14):2826-2835. doi:10.1182/blood-2012-06-435669
 31. Falini B, Mecucci C, Tiacci E, et al; GIMEMA Acute Leukemia Working Party. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med*. 2005;352(3):254-266. doi:10.1056/NEJMoa041974
 32. Döhner K, Schlenk RF, Habdank M, et al. Mutant nucleophosmin (*NPM1*) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood*. 2005;106(12):3740-3746. doi:10.1182/blood-2005-05-2164
 33. Ivey A, Hills RK, Simpson MA, et al; UK National Cancer Research Institute AML Working Group. Assessment of minimal residual disease in standard-risk AML. *N Engl J Med*. 2016;374(5):422-433. doi:10.1056/NEJMoa1507471
 34. Bacher U, Haferlach C, Kern W, Haferlach T, Schnittger S. Prognostic relevance of *FLT3*-TKD mutations in AML: the combination matters—an analysis of 3082 patients. *Blood*. 2008;111(5):2527-2537. doi:10.1182/blood-2007-05-091215
 35. Pratz KW, Lewis M. How I treat *FLT3*-mutated AML. *Blood*. 2017;129(5):565-571. doi:10.1182/blood-2016-09-693648
 36. Stone RM, Mandrekar SJ, Sanford BL, et al. Midostaurin plus chemotherapy for acute myeloid leukemia with a *FLT3* mutation. *N Engl J Med*. 2017;377(5):454-464. doi:10.1056/NEJMoa1614359
 37. Erba HP, Montesinos P, Kim HJ, et al; QuANTUM-First Study Group. Quizartinib plus chemotherapy in newly diagnosed patients with *FLT3*-internal-tandem-duplication-positive acute myeloid leukaemia (QuANTUM-First): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2023;401(10388):1571-1583. doi:10.1016/S0140-6736(23)00464-6
 38. Quizartinib prolonged survival vs placebo plus intensive induction and consolidation therapy followed by single-agent continuation in patients aged 18-75 years with newly diagnosed *FLT3*-ITD+ AML. HemaSphere; 2022; Session Abstract Book for the 27th Congress of the European Hematology Association. Accessed January 12, 2024. https://journals.lww.com/hemasphere/Fulltext/2022/06003/Abstract_Book_for_the_27th_Congress_of_the.1.aspx
 39. Killock D. Newly diagnosed AML: quizartinib improves OS. *Nat Rev Clin Oncol*. 2023;20(8):504. doi:10.1038/s41571-023-00787-6
 40. Lancet JE, Uy GL, Cortes JE, et al. CPX-351 (cytarabine and daunorubicin) liposome for injection versus conventional cytarabine plus daunorubicin in older patients with newly diagnosed secondary acute myeloid leukemia. *J Clin Oncol*. 2018;36(26):2684-2692. doi:10.1200/JCO.2017.77.6112
 41. Jen EY, Wang X, Li M, et al. FDA approval summary: oral azacitidine for continued treatment

- of adults with acute myeloid leukemia unable to complete intensive curative therapy. *Clin Cancer Res*. 2022;28(14):2989-2993. doi:10.1158/1078-0432.CCR-21-4525
42. Wei AH, Döhner H, Pocock C, et al; QUAZAR AML-001 Trial Investigators. Oral azacitidine maintenance therapy for acute myeloid leukemia in first remission. *N Engl J Med*. 2020;383(26):2526-2537. doi:10.1056/NEJMoa2004444
 43. Garcia-Manero G, Gore SD, Cogle C, et al. Phase I study of oral azacitidine in myelodysplastic syndromes, chronic myelomonocytic leukemia, and acute myeloid leukemia. *J Clin Oncol*. 2011;29(18):2521-2527. doi:10.1200/JCO.2010.34.4226
 44. Laille E, Shi T, Garcia-Manero G, et al. Pharmacokinetics and pharmacodynamics with extended dosing of CC-486 in patients with hematologic malignancies. *PLoS One*. 2015;10(8):e0135520. doi:10.1371/journal.pone.0135520
 45. Sekeres MA, Guyatt G, Abel G, et al. American Society of Hematology 2020 guidelines for treating newly diagnosed acute myeloid leukemia in older adults. *Blood Adv*. 2020;4(15):3528-3549. doi:10.1182/bloodadvances.2020001920
 46. Kantarjian H, Ravandi F, O'Brien S, et al. Intensive chemotherapy does not benefit most older patients (age 70 years or older) with acute myeloid leukemia. *Blood*. 2010;116(22):4422-4429. doi:10.1182/blood-2010-03-276485
 47. Zeidan AM, Podoltsev NA, Wang X, et al. Temporal patterns and predictors of receiving no active treatment among older patients with acute myeloid leukemia in the United States: a population-level analysis. *Cancer*. 2019;125(23):4241-4251. doi:10.1002/cncr.32439
 48. DiNardo CD, Wei AH. How I treat acute myeloid leukemia in the era of new drugs. *Blood*. 2020;135(2):85-96. doi:10.1182/blood.2019001239
 49. Konopleva M, Thirman MJ, Pratz KW, et al. Impact of FLT3 mutation on outcomes after venetoclax and azacitidine for patients with treatment-naïve acute myeloid leukemia. *Clin Cancer Res*. 2022;28(13):2744-2752. doi:10.1158/1078-0432.CCR-21-3405
 50. Pollyea DA, DiNardo CD, Arellano ML, et al. Impact of venetoclax and azacitidine in treatment-naïve patients with acute myeloid leukemia and IDH1/2 mutations. *Clin Cancer Res*. 2022;28(13):2753-2761. doi:10.1158/1078-0432.CCR-21-3467
 51. Bernard E, Nannay Y, Hasserjian RP, et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat Med*. 2020;26(10):1549-1556. doi:10.1038/s41591-020-1008-z
 52. Pollyea DA, Pratz KW, Wei AH, et al. Outcomes in patients with poor-risk cytogenetics with or without TP53 mutations treated with venetoclax and azacitidine. *Clin Cancer Res*. 2022;28(24):5272-5279. doi:10.1158/1078-0432.CCR-22-1183
 53. Tiong IS, Hiwase DK, Abro E, et al. Targeting molecular measurable residual disease and low-blast relapse in AML with venetoclax and low-dose cytarabine: a prospective phase II study (VALDAC). *J Clin Oncol*. 2024;JCO2301599. doi:10.1200/JCO.23.01599
 54. Gutman JA, Winters A, Amaya ML, et al. Venetoclax and azacitidine for newly diagnosed non-elderly adult patients with acute myeloid leukemia and adverse risk features. *Blood*. 2020;136(suppl 1):9. doi:10.1182/blood-2020-136120
 55. Short NJ, Daver N, Dinardo CD, et al. Azacitidine, venetoclax, and gilteritinib in newly diagnosed and relapsed or refractory FLT3-mutated AML. *J Clin Oncol*. 2024;42(13):1499-1508. doi:10.1200/JCO.23.01911
 56. Liu Q, Welkie RL, Huang Y, et al. Beat AML S8 Group 2: gilteritinib (GILT) in combination with decitabine (DEC) and venetoclax (VEN) in untreated FLT3 mutated acute myeloid leukemia (AML) patients age ≥60 with high and low variant allele frequency (VAF). *Blood*. 2023;142(suppl 1):5933. doi:10.1182/blood-2023-179609
 57. Montesinos P, Recher C, Vives S, et al. Ivosidenib and azacitidine in IDH1-mutated acute myeloid leukemia. *N Engl J Med*. 2022;386(16):1519-1531. doi:10.1056/NEJMoa2117344
 58. Ozga MP, Dvorak-Kornaus KM, Zhao Q, et al. I-DATA Study: randomized, sequential, open-label study to evaluate the efficacy of IDH targeted/non-targeted versus non-targeted/IDH-targeted approaches in the treatment of newly diagnosed IDH mutated adult AML patients not candidates for intensive induction therapy. *Blood*. 2023;142(suppl 1):1534. doi:10.1182/blood-2023-187060
 59. Ganzel C, Sun Z, Cripe LD, et al. Very poor long-term survival in past and more recent studies for relapsed AML patients: the ECOG-ACRIN experience. *Am J Hematol*. 2018;93(8):1074-1081. doi:10.1002/ajh.25162
 60. DiNardo CD, Maiti A, Rausch CR, et al. 10-Day decitabine with venetoclax for newly diagnosed intensive chemotherapy ineligible, and relapsed or refractory acute myeloid leukaemia: a single-centre, phase 2 trial. *Lancet Haematol*. 2020;7(10):e724-e736. doi:10.1016/S2352-3026(20)30210-6
 61. Stelljes M, Middeke JM, Bug G, et al. In patients with relapsed/refractory AML sequential conditioning and immediate allogeneic stem cell transplantation (allo-HCT) results in similar overall and leukemia-free survival compared to intensive remission induction chemotherapy followed by Allo-HCT: results from the randomized phase III ASAP trial. *Blood*. 2022;140(suppl 1):9-11. doi:10.1182/blood-2022-159962
 62. Stein EM, DiNardo CD, Pollyea DA, et al. Enasidenib in mutant IDH2 relapsed or refractory acute myeloid leukemia. *Blood*. 2017;130(6):722-731. doi:10.1182/blood-2017-04-779405
 63. de Botton S, Montesinos P, Schuh AC, et al. Enasidenib vs conventional care in older patients with late-stage mutant-IDH2 relapsed/refractory AML: a randomized phase 3 trial. *Blood*. 2023;141(2):156-167. doi:10.1182/blood.2021014901
 64. Venugopal S, Watts J. Olutasidenib: from bench to bedside. *Blood Adv*. 2023;7(16):4358-4365. doi:10.1182/bloodadvances.2023009854
 65. Popovici-Muller J, Lemieux RM, Artin E, et al. Discovery of AG-120 (ivosidenib): a first-in-class mutant IDH1 inhibitor for the treatment of IDH1 mutant cancers. *ACS Med Chem Lett*. 2018;9(4):300-305. doi:10.1021/acsmedchemlett.7b00421
 66. Caravella JA, Lin J, Diebold RB, et al. Structure-based design and identification of FT-2102 (olutasidenib), a potent mutant-selective IDH1 inhibitor. *J Med Chem*. 2020;63(4):1612-1623. doi:10.1021/acs.jmedchem.9b01423
 67. DiNardo CD, Stein EM, de Botton S, et al. Durable remissions with ivosidenib in IDH1-mutated relapsed or refractory AML. *N Engl J Med*. 2018;378(25):2386-2398. doi:10.1056/NEJMoa1716984
 68. de Botton S, Fenaux P, Yee K, et al. Olutasidenib (FT-2102) induces durable complete remissions in patients with relapsed or refractory IDH1-mutated AML. *Blood Adv*. 2023;7(13):3117-3127. doi:10.1182/bloodadvances.2022009411
 69. Levis M, Perl AE. Gilteritinib: potent targeting of FLT3 mutations in AML. *Blood Adv*. 2020;4(6):1178-1191. doi:10.1182/bloodadvances.2019000174
 70. Perl AE, Martinelli G, Cortes JE, et al. Gilteritinib or chemotherapy for relapsed or refractory FLT3-mutated AML. *N Engl J Med*. 2019;381(18):1728-1740. doi:10.1056/NEJMoa1902688
 71. Smith CC, Levis MJ, Perl AE, Hill JE, Rosales M, Bahceci E. Molecular profile of FLT3-mutated relapsed/refractory patients with AML in the phase 3 ADMIRAL study of gilteritinib. *Blood Adv*. 2022;6(7):2144-2155. doi:10.1182/bloodadvances.2021006489
 72. Issa GC, Aldoss I, DiPersio J, et al. The menin inhibitor revumenib in KMT2A-rearranged or NPM1-mutant leukaemia. *Nature*. 2023;615(7954):920-924. doi:10.1038/s41586-023-05812-3
 73. DiNardo CD, Tiong IS, Quaglieri A, et al. Molecular patterns of response and treatment failure after frontline venetoclax combinations in older patients with AML. *Blood*. 2020;135(11):791-803. doi:10.1182/blood.2019003988
 74. Perner F, Stein EM, Wenge DV, et al. MEN1 mutations mediate clinical resistance to menin inhibition. *Nature*. 2023;615(7954):913-919. doi:10.1038/s41586-023-05755-9
 75. Venugopal S, Watts J. The future paradigm of HMA + VEN or targeted inhibitor approaches: sequencing or triplet combinations in AML therapy. *Hematology Am Soc Hematol Educ Program*. 2023;2023(1):192-197. doi:10.1182/hematology.2023000429
 76. Madarang E, Lykon J, Zhao W, et al. Toxicity and outcomes in octo- and nonagenarians with AML treated with venetoclax and hypomethylating agent therapy. *Blood*. 2022;140(suppl 1):3306-3308. doi:10.1182/blood-2022-170771
 77. Garcia S, Bertoli S, Sallman DA, et al. Acute myeloid leukemia patients who stopped venetoclax or/and azacitidine for other reasons than progression have a prolonged treatment free remission and overall survival: a Filo study. *Blood*. 2023;142(suppl 1):161. doi:10.1182/blood-2023-185437
 78. Twain M. *The Innocents Abroad*. American Publishing Co; 1869.